

February "Fast Facts About Fast Track Anesthesia" References

- White PF. Ambulatory anesthesia advances into the new millennium. *Anesth Analg* 2000;90:1234-5
- Gupta A, *et al.* Comparison of recovery profile after ambulatory anesthesia with propofol, isoflurane, sevoflurane and desflurane: a systemic review. *Anesth Analg* 2004;98(3):632-41
- Tang J, Chen L, White PF, *et al.* Fast-track office-based anesthesia: a comparison of propofol versus desflurane with antiemetic prophylaxis in spontaneously breathing patients. *Anesth Analg* 2001;92:95-9
- Coloma M, Zhou T, White PF, *et al.* Fast-tracking after outpatient laparoscopy: reasons for failure after propofol, sevoflurane, and desflurane anesthesia. *Anesth Analg* 2001;93:112-5
- Hadzic A, Arliss J, Kerimoglu B, *et al.* A comparison of infraclavicular nerve block versus general anesthesia for hand and wrist day-case surgeries. *Anesthesiology* 2004;101:127-32
- Aldrete JA. The post-anesthesia score revisited. *J Clin Anesth* 1995;7:89-91
- Marshall SI, *et al.* Discharge criteria and complication after ambulatory surgery. *Anesth Analg* 1999;88:508-16
- White PF, Song D. New criteria for fast-tracking after outpatient anesthesia: a comparison with the modified Aldrete's scoring system. *Anesth Analg* 1999;88:1069-72
- Watkins AC, White PF. Fast-tracking after ambulatory surgery. *J Perianesth Nurs* 2001;16:379-87
- Macario A, Vitez TS, Dunn S, *et al.* Where are the costs in perioperative care? Analysis of hospital costs and charges for inpatient surgical care. *Anesthesiology* 1995;82:94-101
- Dexter F, Tinker JH. Analysis of strategies to decrease postanesthesia care unit costs. *Anesthesiology* 1995;82:94-101
- Dexter F, Macario A, Manberg PJ, *et al.* Computer simulation to determine how rapid anesthetic recovery protocols to decrease the time of emergence or increase phase I postanesthesia care unit bypass rate affect staffing of an ambulatory surgery center. *Anesth Analg* 1999;88:842-8
- White PF, Ma H, Tang J, *et al.* Does the use of electroencephalographic bispectral index or auditory evoked potential index monitoring facilitate recovery after desflurane anesthesia in the ambulatory setting? *Anesthesiology* 2004;100(4):811-7
- Pavlin DJ, *et al.* The effect of bispectral index monitoring on end-tidal gas concentration and recovery duration after outpatient anesthesia. *Anesth Analg* 2001;93:613-9
- Kalkman CJ, *et al.* Monitors of depth of anesthesia, *quo vadis?* *Anesthesiology* 2002;96:784-7
- Ahmad S, *et al.* Impact of bispectral index monitoring on fast-tracking of gynecologic patients undergoing laparoscopic surgery. *Anesthesiology* 2003;98:849-52

- Michaloliakou C, Chung F, Sharma S. Preoperative multimodal analgesia facilitates recovery after ambulatory laparoscopic cholecystectomy. *Anesth Analg* 1996; 82:44-51
- Joshi GP. Pain management after ambulatory surgery. *Ambulatory Surgery* 1999;7:3-12
- Scuderi PE, James RL, Harris L, *et al.* Multimodal antiemetic management prevents early postoperative vomiting after outpatient laparoscopy. *Anesth Analg* 2000;91:1408-14
- White PF. The role of non-opioid analgesic techniques in the management of pain after ambulatory surgery. *Anesth Analg* 2002;94:577-85
- Ma H, *et al.* Perioperative rofecoxib improves early recovery after outpatient herniorrhaphy. *Anesth Analg* 2004;98:970-5
- Recart A, *et al.* The efficacy of celecoxib premedication on postoperative pain and recovery times after ambulatory surgery: a dose-ranging study. *Anesth Analg* 2003;96:1631-5
- Joshi GP, *et al.* Effective treatment of laparoscopic cholecystectomy pain with intravenous followed by oral COX-2 specific inhibitor. *Anesth Analg* 2003;98(2):336-42
- Apfel C, Kortilla K, Abdalla M, *et al.* A factorial trial of six interventions for the prevention of postoperative nausea and vomiting. *NEJM* 2004;350:2441-2451
- White PF. Prevention of postoperative nausea and vomiting - a multimodal solution to a persistent problem (Editorial). *NEJM* 2004;350(24):2511-2
- Scuderi PE, *et al.* Multimodal antiemetic management prevents early postoperative vomiting after outpatient laparoscopy. *Anesth Analg* 2000;91:1408-14
- Yogendran S, Asokumar B, Cheng D, *et al.* A prospective, randomized double-blind study of the effect of intravenous fluid therapy on adverse outcomes after outpatient surgery. *Anesth Analg* 1995;80:682-6