

**March “How to Make Regional Anesthesia Work” References:**

1. Parnass SM, McCarthy RJ, Back B. et al. A prospective evaluation of epidural versus general anesthesia for outpatient arthroscopy. *Anesthesiology* 1980; 73: A23.
2. Neal JM, Deck JJ, Lewis MA et al. A double blind comparison of epidural 2-chloroprocaine versus lidocaine for outpatient knee arthroscopy. *Anesthesiology* 1993;79:A12
3. Kopacz DJ, Mulroy MF. Chlorprocaine and lidocaine decrease hospital stay and admission rate after outpatient epidural anesthesia. *Reg Anesthesia* 1990; 15:19-25.
4. Rodgers A, Walker N, Schug SA, et al. Regional anaesthesia reduces postoperative mortality and morbidity: results from an overview of randomized trials. *Lancet* 1999 (submitted for publication).
5. Greenberg CP, Wald A, Roth JM. Cost conscious anesthetic choices for ambulatory surgery. *Anesthesiology* 1994; 81: A1202
6. Allen HW, Mulroy MF, Fundis K, et al. Regional anesthesia versus propofol general anesthesia for outpatient hand surgery. *Anesthesiology* 1993;79:A1
7. Hadzic A, Vloka JD, Kuroda MM, et al. The Practice of Peripheral Nerve Blocks in the United States: a national survey. *Reg Anesth Pain Med* 1998, 241-6.
8. Liu S, Carpenter RL, Neal JM. Epidural Anesthesia and analgesia. Their role in postoperative outcome. *Anesthesiology* 1995;82:1474-506.
9. Tanaka K, Watanabe R, Harada T et al. Extensive application of epidural anesthesia and analgesia in a University Hospital: Incidence of Complications Related to Technique. *Reg Anes and Pain Med* 1993; 18;34-8.
10. Bromage P. Epidural Analgesia. Philadelphia. WB Saunders, 1978, p. 197.
11. Pollock JE, Neal JM, Stephenson CA, et al. Prospective study of the incidence of transient radicular irritation in patients undergoing spinal anesthesia. *Anesthesiology* 1996, 84:1361-7.
12. Freedman JM, Li DK, Drasner K, et al. Transient neurologic symptoms after spinal anesthesia. *Anesthesiology* 1998; 89: 633-41
13. Ben-David B, Maryanovsky M, Gurevitch A, et al. A Comparison of Minidose Lidocaine-Fentanyl and Conventional Dose Lidocaine Spinal Anesthesia. *Anesth Analg* 2000; 91:865-70
14. Goldberg ME, Gregg C, Lorijani GE, et al. A comparison of three methods of axillary approach to brachial plexus for upper extremity surgery. *Anesthesiology* 1987; 66:814-6
15. Urmey WF, Talts KH, Sharrock NE. One hundred percent incidence of hemidiaphragmatic paresis associated with intrascalene brachial plexus anesthesia as diagnosed by ultrasonography.
16. Allen HW, Liu SS, Ware PD, et al. Peripheral nerve block improves analgesia after total knee replacement surgery. *Anesthesia & Analgesia* 1998; 87:93-7.