

## References

1. Drummond JC. Monitoring Depth of Anesthesia. *Anesthesiology* 2000; 93(3)
2. Stanski DR, Shafer SL. Measuring depth of anesthesia. *Anesthesia, Sixth Edition*. Edited by Miller RD. United States, Elsevier Churchill Livingstone, 2005, Chap 31, pp 1227-64.
3. Bruhn J, Myles PS, et al. Depth of anaesthesia monitoring: what's available, what's validated and what's next? *Br J Anaesth*. 2006; 97:85-94.
4. Jensen EW, Litvan H, et al. Pitfalls and challenges when assessing the depth of hypnosis during general anaesthesia by clinical signs and electronic indices. *Acta Anaesthesiol Scand*. 2004; 48:1260-67.
5. Glass PS, Bloom M, Kears L, et al. Bispectral analysis measures sedation and memory effects of propofol, midazolam, isoflurane, and alfentanil in healthy volunteers. *Anesthesiology*. 1997; 86(4):836-47.
6. Mahla M, Black S, Cucchiara RF. Neurologic Monitoring. *Anesthesia, Sixth edition*, Edited by Miller RD. United States, Elsevier Churchill Livingstone, 2005, Chap 38, pp 1516-17
7. Sneyd JR. Editorial III: Remembering awareness. *Br J of Anaesth*. 2004; 93(3):324-6.
8. Sebel PS, Bowdle TA, Ghoneim MM, et al. The incidence of awareness during anesthesia: A Multi-center United States Study. *Anesth and Analg*. 2004; 99:833-9
9. Osterman JE, Hopper J, Heran WJ, et al. Awareness under anesthesia and the development of posttraumatic stress disorder. *Gen Hosp Psychiatry*. 2001; 23:198-204.